

Equivariant Operators between some Modules of the Lie Algebra of Vector Fields*

N. Poncin

February 1, 2008

Abstract

The space $\mathcal{D}_p^k$ of differential operators of order $\leq k$, from the differential forms of degree p of a smooth manifold M into the functions of M , is a module over the Lie algebra of vector fields of M , when it's equipped with the natural Lie derivative. In this paper, we compute all equivariant i.e. intertwining operators $T : \mathcal{D}_p^k \rightarrow \mathcal{D}_q^\ell$ and conclude that the preceding modules of differential operators are never isomorphic. We also answer a question of P. Lecomte, who observed that the restriction of some homotopy operator—introduced in [2]—to $\mathcal{D}_p^k$ is equivariant for small values of k and p .

Mathematics Subject Classification (2000): 17B66, 16D99, 58J99.

Keywords: Lie algebra of vector fields, modules of differential operators, intertwining operators.

1 Introduction

Let M be a smooth, Hausdorff, second countable, connected manifold of dimension m .

Denote by $\Omega^p(M)$ the space of differential forms of degree p of M , by N the space $C^\infty(M)$ of smooth functions of M , and by $\mathcal{D}_p^k$ the space of differential operators of order smaller than or equal to k , from $\Omega^p(M)$ into N . If $X \in Vect(M)$, where $Vect(M)$ is the Lie algebra of vector fields of M , and $D \in \mathcal{D}_p^k$, the Lie derivative

$$L_X D = L_X \circ D - D \circ L_X \quad (1)$$

is a differential operator of order at most k and so $(\mathcal{D}_p^k, L)$ is a $Vect(M)$ -module.

In this paper, we shall determine all the spaces $\mathcal{I}_{p,q}^{k,\ell}$ of *equivariant* operators from $\mathcal{D}_p^k$ into $\mathcal{D}_q^\ell$, that is all operators $T : \mathcal{D}_p^k \rightarrow \mathcal{D}_q^\ell$ such that

$$L_X \circ T = T \circ L_X, \forall X \in Vect(M). \quad (2)$$

In [3], P. Lecomte, P. Mathonet, and E. Tousset computed all linear equivariant mappings between modules of differential operators acting on densities. We

*This work was supported by MCESR Grant MEN/CUL/99/007. The author thanks P. Lecomte and P. Mathonet for helpful comments.

solve here an analogous problem, thus answering a question of P. Lecomte, who noticed that some homotopy operator, which locally coincides—up to a coefficient—with the Koszul differential (see [2]), is equivariant if it's restricted to low order differential operators and asked whether this property still holds for higher orders.

2 Local representation

The formalism described in this section was also used in [5] and [6].

Consider an open subset U of $\mathbb{R}^m$, two real finite-dimensional vector spaces E and F , and some local operator

$$O \in \mathcal{L}(C^\infty(U, E), C^\infty(U, F))_{loc}.$$

The operator is fully defined by its values on the products fe , $f \in C^\infty(U)$, $e \in E$. A well known theorem of J. Peetre (see [4]) states that it has the form

$$O(fe) = \sum_{\alpha} O_{\alpha}(\partial^{\alpha}(fe)) = \sum_{\alpha} O_{\alpha}(e)\partial^{\alpha}f,$$

where $\partial_x^{\alpha} = \partial_{x^1}^{\alpha_1} \dots \partial_{x^m}^{\alpha_m}$ and $O_{\alpha} \in C^\infty(U, \mathcal{L}(E, F))$. Moreover, the coefficients O_{α} are well determined by O and the series is locally finite (it is finite, if U is relatively compact).

We shall symbolize the partial derivative $\partial^{\alpha}f$ by the monomial $\xi^{\alpha} = \xi_1^{\alpha_1} \dots \xi_m^{\alpha_m}$ in the components $\xi_1, \dots, \xi_m$ of some linear form $\xi \in (\mathbb{R}^m)^*$, or—at least mentally—even by $\xi^{\alpha}f$, if this is necessary to avoid confusion. The operator O is thus represented by the polynomial

$$\mathcal{O}(\xi; e) = \sum_{\alpha} O_{\alpha}(e)\xi^{\alpha}.$$

When identifying the space $Pol(\mathbb{R}^m)^*$ of polynomials on $(\mathbb{R}^m)^*$ with the space $\vee \mathbb{R}^m$ of symmetric contravariant tensors of $\mathbb{R}^m$, one has $\mathcal{O} \in C^\infty(U, \vee \mathbb{R}^m \otimes \mathcal{L}(E, F))$. Let's emphasize that the form ξ symbolizes the derivatives in O that act on the argument $fe \in C^\infty(U, E)$, while $e \in E$ represents this argument. In the sequel, we shall no longer use different notations for the operator O and its representative polynomial $\mathcal{O}$; in order to simplify notations, it's helpful to use even the same typographical sign, when referring to the argument fe and its representation e .

Let's for instance look for the local representation of (1). If $D \in \mathcal{D}_p^k$, its restriction $D|_U$ (or simply D , if no confusion is possible) to a domain U of local coordinates of M , is a local operator from $C^\infty(U, \wedge^p(\mathbb{R}^m)^*)$ into $C^\infty(U)$ that is represented by $D(\omega) \simeq D(\xi; \omega)$, where $\omega \in C^\infty(U, \wedge^p(\mathbb{R}^m)^*)$ on the l.h.s. and $\omega \in \wedge^p(\mathbb{R}^m)^*$ on the r.h.s., and where ξ represents the derivatives acting on ω . The Lie derivative of $D(\omega)$ with respect to a vector field $X \in C^\infty(U; \mathbb{R}^m)$, is then represented by $L_X(D(\omega)) \simeq \langle X, \eta + \xi \rangle D(\xi; \omega)$. Here $\eta \in (\mathbb{R}^m)^*$ is associated to D and $\langle X, \eta + \xi \rangle$ denotes the evaluation of $X \in \mathbb{R}^m$ on $\eta + \xi$. When associating ζ to X , one gets $D(L_X\omega) \simeq D(\xi + \zeta; \langle X, \xi \rangle \omega + \zeta \wedge i_X\omega)$ and

$$(L_X D)(\omega) \simeq \langle X, \eta \rangle D(\xi; \omega) - \langle X, \xi \rangle \tau_{\zeta} D(\xi; \omega) - D(\xi + \zeta; \zeta \wedge i_X\omega), \quad (3)$$

where $\tau_{\zeta} D(\xi; \omega) = D(\xi + \zeta; \omega) - D(\xi; \omega)$.

3 Locality of equivariant operators

Let U be a domain of local coordinates of M . If $D \in \mathcal{D}_p^k|_U$ ($\mathcal{D}_p^k|_U$ is defined similarly to $\mathcal{D}_p^k$, but for $M = U$), its representation is a polynomial D in $C^\infty(U, E_p^k)$, where $E_p^k = \vee^{\leq k} \mathbb{R}^m \otimes \wedge^p \mathbb{R}^m$, or—equivalently—in the space of smooth sections $\Gamma(\vee^{\leq k} TU \otimes \wedge^p TU)$. We denote by $\sigma(D)$ or simply σ the homogeneous part of maximal degree of this polynomial i.e. the principal symbol of the considered differential operator. If L^t is the natural Lie derivative of tensor fields, one easily verifies that

$$(L_X^t(\sigma(D))) (\xi; \omega) = \langle X, \eta \rangle \sigma(\xi; \omega) - \langle X, \xi \rangle (\zeta \partial_\xi) \sigma(\xi; \omega) - \sigma(\xi; \zeta \wedge i_X \omega),$$

where η and ζ are associated to σ and X respectively, and where $\zeta \partial_\xi$ denotes the derivation with respect to ξ in the direction of ζ . If L^{op} is the formerly defined Lie derivative L of differential operators, it follows from (3) that $(\sigma(L_X^{op} D)) (\xi; \omega) = (L_X^t(\sigma(D))) (\xi; \omega)$ i.e. that the principal symbol is equivariant with respect to all vector fields.

We are now prepared to prove the following lemma:

Lemma 1 *Every equivariant operator $T \in \mathcal{I}_{p,q}^{k,\ell}$ is local.*

Proof. It suffices to show that the family $\mathcal{L}_p^k = \{L_X : \mathcal{D}_p^k \rightarrow \mathcal{D}_p^k \mid X \in \text{Vect}(M)\}$ is closed with respect to locally finite sums and is locally transitive (lt); this means that each point of M has some neighborhood Ω , such that for every open subset $\omega \subset \Omega$ and every $D \in \mathcal{D}_p^k$ with compact support in ω , D can be decomposed into Lie derivatives,

$$D = \sum_{i=1}^n L_{X_i} D_i \quad (D_i \in \mathcal{D}_p^k, \text{supp } X_i \subset \omega, \text{supp } D_i \subset \omega), \quad (4)$$

with n independent of D and Ω . Indeed, proposition 3 of [1] states that $\mathcal{L}_p^k$ is then globally transitive i.e. that (4) holds for every open subset ω of M and every $D \in \mathcal{D}_p^k$ with support in ω . Take now a differential operator $D \in \mathcal{D}_p^k$ that vanishes in an open subset V of M and let x_0 be an arbitrary point of V . Since $\text{supp } D \subset \omega = M \setminus \omega_0$ for some neighborhood ω_0 of x_0 , D has form (4) and $T(D)(x_0) = \sum_{i=1}^n (L_{X_i}(T(D_i)))(x_0) = 0$.

In order to confirm local transitivity (LT), let's point out that example 12 of [1] shows that the family $\mathcal{L}_s^r = \{L_X : \Gamma(\otimes_s^r M) \rightarrow \Gamma(\otimes_s^r M) \mid X \in \text{Vect}(M)\}$ is lt, if

$$s - r \neq m. \quad (5)$$

Moreover, the domains U of the charts of M that correspond to cubes in $\mathbb{R}^m$, play the role of the neighborhoods Ω in the definition of LT.

LT of $\mathcal{L}_p^k$ may now be proved as follows. If $x \in M$, take $\Omega = U$ and if $D \in \mathcal{D}_p^k$ is a differential operator with compact support in some open subset ω of U , note that its principal symbol $\sigma(D)$ is a contravariant tensor field with compact support in ω . Hence condition (5) is satisfied and

$$\sigma(D) = \sum_{i=1}^n L_{X_i}^t \sigma(D_i) = \sigma \left(\sum_{i=1}^n L_{X_i}^{op} D_i \right)$$

($D_i \in \mathcal{D}_p^k$; $\text{supp } X_i, \text{supp } D_i \subset \omega$; n independent of D and U). Thus $D - \sum_{i=1}^n L_{X_i}^{op} D_i \in \mathcal{D}_p^{k-1}$ and we conclude by iteration. ■

4 Local expression of the equivariance equation

Let U be a connected, relatively compact domain of local coordinates of M .

Recall that if $D \in \mathcal{D}_p^k|_U$, its representation is a polynomial $D \in C^\infty(U, E_p^k)$, where $E_p^k = \vee^{\leq k} \mathbb{R}^m \otimes \wedge^p \mathbb{R}^m$.

We identify $\mathcal{D}_p^k|_U$ with $C^\infty(U, E_p^k)$. Thus, if $T \in \mathcal{I}_{p,q}^{k,\ell}$, the restriction $T|_U$ is a local operator from $C^\infty(U, E_p^k)$ into $C^\infty(U, E_q^\ell)$, with representation $T(\eta; D)(\xi; \omega)$ ($\eta, \xi \in (\mathbb{R}^m)^*, D \in E_p^k, \omega \in \wedge^q(\mathbb{R}^m)^*$).

It's easily checked that equivariance condition (2) locally reads

$$\begin{aligned} (X.T)(\eta; D)(\xi; \omega) - \langle X, \eta \rangle (\tau_\zeta T(\eta; D))(\xi; \omega) \\ - \langle X, \xi \rangle \tau_\zeta (T(\eta; D))(\xi; \omega) + T(\eta + \zeta; X \tau_\zeta D)(\xi; \omega) \\ - T(\eta; D)(\xi + \zeta; \zeta \wedge i_X \omega) + T(\eta + \zeta; D(\cdot + \zeta; \zeta \wedge i_X \cdot))(\xi; \omega) = 0, \end{aligned} \quad (6)$$

where $X.T$ is obtained by derivation of the coefficients of T in the direction of X .

Take in equation (6) the terms of degree 0 in ζ :

$$(X.T)(\eta; D)(\xi; \omega) = 0.$$

This means that the coefficients of T are constant.

The terms of degree 1 lead to the equation

$$\begin{aligned} \langle X, \eta \rangle (\zeta \partial_\eta) T(\eta; D)(\xi; \omega) \\ - T(\eta; X(\zeta \partial_\xi) D)(\xi; \omega) - T(\eta; D(\cdot; \zeta \wedge i_X \cdot))(\xi; \omega) \\ + \langle X, \xi \rangle (\zeta \partial_\xi) T(\eta; D)(\xi; \omega) + T(\eta; D)(\xi; \zeta \wedge i_X \omega) = 0, \end{aligned}$$

which, if ρ denotes the natural action of $gl(m, \mathbb{R})$, may be rewritten

$$\rho(X \otimes \zeta) (T(\eta; D)(\xi; \omega)) = 0.$$

Note that $T(\eta; D)(\xi; \omega)$ is completely characterized by $T(\eta; Y^r \otimes (X_1 \wedge \dots \wedge X_p))(\xi; \nu^1 \wedge \dots \wedge \nu^q)$ ($Y, X_i \in \mathbb{R}^m, \nu^j \in (\mathbb{R}^m)^*, r \in \{0, \dots, k\}$). This last expression is a polynomial in Y, X_i and in η, ξ, ν^j . It thus follows from the description of invariant polynomials under the action of $gl(m, \mathbb{R})$ (see [7]), that it is a polynomial $\mathcal{T}_r(\langle Y, \xi \rangle, \langle Y, \eta \rangle, \langle Y, \nu^j \rangle, \langle X_i, \xi \rangle, \langle X_i, \eta \rangle, \langle X_i, \nu^j \rangle)$ in the evaluations of the vectors on the linear forms.

In order to determine the most general structure of $\mathcal{T}_r$, observe that this polynomial is homogeneous of degree r in Y and degree 1 in the X_i 's and the ν^j 's, and that furthermore it's antisymmetric in the X_i 's and the ν^j 's. It follows from the skew-symmetry in the ν^j 's, that Y is evaluated on at most one ν^j , so that $q \leq p + 1$, and from the skew-symmetry in the X_i 's, that ξ and η are evaluated on at most one X_i , so that $q \geq p - 2$. Finally, q is $p - 2, p - 1, p$ or $p + 1$. Set now $\Lambda = X_1 \wedge \dots \wedge X_p, \omega = \nu^1 \wedge \dots \wedge \nu^q, u = \langle Y, \xi \rangle$ and $v = \langle Y, \eta \rangle$. The following possible forms of the terms of $\mathcal{T}_r$ and the corresponding conditions on

p and q are immediate consequences of the preceding observations.

term type	condition	term form
(1) no $\langle Y, \nu^j \rangle$		
(1.1) no $\langle X_i, \xi \rangle$	no $\langle X_i, \eta \rangle$ $q = p$	$v^s u^{r-s} \langle \Lambda, \omega \rangle$
(1.2) one $\langle X_i, \xi \rangle$	no $\langle X_i, \eta \rangle$ $q = p - 1$	$v^s u^{r-s} \langle i_\xi \Lambda, \omega \rangle$
(1.3) no $\langle X_i, \xi \rangle$	one $\langle X_i, \eta \rangle$ $q = p - 1$	$v^s u^{r-s} \langle i_\eta \Lambda, \omega \rangle$
(1.4) one $\langle X_i, \xi \rangle$	one $\langle X_i, \eta \rangle$ $q = p - 2$	$v^s u^{r-s} \langle i_\xi i_\eta \Lambda, \omega \rangle$
(2) one $\langle Y, \nu^j \rangle$		
(2.1) no $\langle X_i, \xi \rangle$	no $\langle X_i, \eta \rangle$ $q = p + 1$	$v^s u^{r-s-1} \langle Y \wedge \Lambda, \omega \rangle$
(2.2) one $\langle X_i, \xi \rangle$	no $\langle X_i, \eta \rangle$ $q = p$	$v^s u^{r-s-1} \langle Y \wedge i_\xi \Lambda, \omega \rangle$
(2.3) no $\langle X_i, \xi \rangle$	one $\langle X_i, \eta \rangle$ $q = p$	$v^s u^{r-s-1} \langle Y \wedge i_\eta \Lambda, \omega \rangle$
(2.4) one $\langle X_i, \xi \rangle$	one $\langle X_i, \eta \rangle$ $q = p - 1$	$v^s u^{r-s-1} \langle Y \wedge i_\xi i_\eta \Lambda, \omega \rangle$

(7)

Remark 1 These structures of $\mathcal{T}_r$ show that a priori equivariant operators are mappings $T : \mathcal{D}_p^k \longrightarrow \mathcal{D}_{p-2}^{k+1}$, $T : \mathcal{D}_p^k \longrightarrow \mathcal{D}_{p-1}^{k+1}$, $T : \mathcal{D}_p^k \longrightarrow \mathcal{D}_p^k$ or $T : \mathcal{D}_p^k \longrightarrow \mathcal{D}_{p+1}^{k-1}$.

It's interesting to rewrite (6) in terms of the $\mathcal{T}_r$'s ($r \in \{0, \dots, k\}$). To simplify notations, we set

$$\begin{aligned} \lambda &= \langle X, \xi \rangle & \mu &= \langle X, \eta \rangle & \nu &= \langle X, \zeta \rangle & \pi^j &= \langle X, \nu^j \rangle \\ u &= \alpha_0 = \langle Y, \xi \rangle & v &= \beta_0 = \langle Y, \eta \rangle & \gamma_0 &= \langle Y, \zeta \rangle & \delta_0^j &= \langle Y, \nu^j \rangle \\ \alpha_i &= \langle X_i, \xi \rangle & \beta_i &= \langle X_i, \eta \rangle & \gamma_i &= \langle X_i, \zeta \rangle & \delta_i^j &= \langle X_i, \nu^j \rangle \end{aligned}$$

Substitute in (6), $Y^r \otimes (X_1 \wedge \dots \wedge X_p)$ and $\nu^1 \wedge \dots \wedge \nu^q$ to D and ω respectively, and use subscript $\iota \in \{0, \dots, p\}$. Note that:

- Taylor expansion gives

$$T(\eta; D)(\xi + \zeta; \omega) = \sum \frac{1}{a!} (\zeta \partial_\xi)^a T(\eta; D)(\xi; \omega) = \sum \frac{1}{a!} (\zeta \partial_\xi)^a \mathcal{T}_r(\alpha_\iota, \beta_\iota, \delta_\iota^j)$$

- the computation of the successive directional derivatives $(\zeta \partial_\xi)^a$ of the composite function $\mathcal{T}_r(\alpha_\iota, \beta_\iota, \delta_\iota^j)$ leads to the same rule than the computation of the successive powers of a sum of $p + 1$ real terms:

$$\begin{aligned} & (\zeta \partial_\xi)^a \mathcal{T}_r(\alpha_\iota, \beta_\iota, \delta_\iota^j) \\ &= \sum_{\varrho_0 + \dots + \varrho_p = a} \frac{a!}{\varrho_0! \dots \varrho_p!} \gamma_0^{\varrho_0} \dots \gamma_p^{\varrho_p} \left(\partial_{\alpha_0}^{\varrho_0} \dots \partial_{\alpha_p}^{\varrho_p} \mathcal{T}_r \right) (\alpha_\iota, \beta_\iota, \delta_\iota^j) \end{aligned}$$

-

$$\tau_\zeta Y^r = \sum_{a=0}^{r-1} \binom{r}{a} \gamma_0^{r-a} Y^a$$

•

$$XY^a = \frac{1}{a+1} (X \partial_Y) Y^{a+1}$$

•

$$\begin{aligned} & (X_1 \wedge \dots \wedge X_p)(\zeta \wedge i_X \cdot) \\ &= X \wedge i_\zeta (X_1 \wedge \dots \wedge X_p) \\ &= \sum_{b=1}^p \gamma_b (X \partial_{X_b}) (X_1 \wedge \dots \wedge X_p) \end{aligned} \quad (8)$$

Transform now the terms of (6) in conformity with the preceding hints. If $(\partial \mathcal{T})(\cdot) = (\partial \mathcal{T})(\alpha_\iota, \cdot, \delta_\iota^j)$ and if $d_r(\xi)$ denotes the degree of $\mathcal{T}_r$ in ξ , the local equivariance equation (6) finally reads in terms of the $\mathcal{T}_r$'s:

$$\begin{aligned} & \mu(\mathcal{T}_r(\beta_\iota + \gamma_\iota) - \mathcal{T}_r(\beta_\iota)) \\ &+ \lambda \sum_{a=1}^{d_r(\xi)} \sum_{\varrho_0 + \dots + \varrho_p = a} \frac{1}{\varrho_0! \dots \varrho_p!} \gamma_0^{\varrho_0} \dots \gamma_p^{\varrho_p} \left(\partial_{\alpha_0}^{\varrho_0} \dots \partial_{\alpha_p}^{\varrho_p} \mathcal{T}_r \right) (\beta_\iota) \\ &+ \sum_{b=1}^q \pi^b \sum_{a=0}^{d_r(\xi)} \sum_{\iota=0}^p \gamma_\iota \\ & \sum_{\varrho_0 + \dots + \varrho_p = a} \frac{1}{\varrho_0! \dots \varrho_p!} \gamma_0^{\varrho_0} \dots \gamma_p^{\varrho_p} \left(\partial_{\alpha_0}^{\varrho_0} \dots \partial_{\alpha_p}^{\varrho_p} \partial_{\delta_\iota^b} \mathcal{T}_r \right) (\beta_\iota) \\ &- \sum_{a=0}^{r-1} \binom{r}{a} \frac{1}{a+1} \gamma_0^{r-a} \left[\lambda (\partial_{\alpha_0} \mathcal{T}_{a+1})(\beta_\iota + \gamma_\iota) \right. \\ & \quad \left. + (\mu + \nu) (\partial_{\beta_0} \mathcal{T}_{a+1})(\beta_\iota + \gamma_\iota) \right. \\ & \quad \left. + \sum_{j=1}^q \pi^j \left(\partial_{\delta_0^j} \mathcal{T}_{a+1} \right) (\beta_\iota + \gamma_\iota) \right] \\ &- \sum_{b=1}^p \gamma_b \sum_{a=0}^r \binom{r}{a} \gamma_0^{r-a} \left[\lambda (\partial_{\alpha_b} \mathcal{T}_a)(\beta_\iota + \gamma_\iota) \right. \\ & \quad \left. + (\mu + \nu) (\partial_{\beta_b} \mathcal{T}_a)(\beta_\iota + \gamma_\iota) \right. \\ & \quad \left. + \sum_{j=1}^q \pi^j \left(\partial_{\delta_b^j} \mathcal{T}_a \right) (\beta_\iota + \gamma_\iota) \right] = 0, \end{aligned} \quad (9)$$

for each $r \in \{0, \dots, k\}$.

Remark 2 *In the sequel, we suppose that dimension m is not only $\geq \sup(p, q)$, but also $\geq \inf\{p+2, q+3\}$. In the skipped borderline cases, results have to be completed, but proofs are not elegant.*

We claim that (9) is a polynomial identity in the independent variables $\lambda, \dots, \delta_p^q$. Indeed, since as well the vectors X, Y, X_i as the forms ξ, η, ζ, ν^j are arbitrary, take $X = e_1, Y = e_2, X_i = e_{i+2}$ (e_k : canonical basis of $\mathbb{R}^m$) and let the $p+2$ first components of the preceding forms in the dual basis of e_k vary in $\mathbb{R}$, if $\inf\{p+2, q+3\} = p+2$; proceed similarly but exchange roles of vectors and forms, if $\inf\{p+2, q+3\} = q+3$.

When seeking in (9) the terms of degree 1 in ν and γ_0 , and of degree 0 in $\gamma_1, \dots, \gamma_p$ (in the sequel we shall denote these terms by $(\nu)^1 \gamma_0^1 \gamma_1^0 \dots \gamma_p^0$), one gets

$$\partial_{\beta_0} \mathcal{T}_r = 0, \quad (10)$$

where $\mathcal{T}_0$ is a priori independent of Y and $\beta_0 = \langle Y, \eta \rangle$. The terms in $(\nu)^1 \gamma_0^0 \gamma_1^0 \dots \gamma_i^1 \dots \gamma_p^0$ ($i \in \{1, \dots, p\}$, $p > 0$), $(\lambda)^1 \gamma_0^2 \gamma_1^0 \dots \gamma_p^0$, $(\lambda)^1 \gamma_0^1 \gamma_1^0 \dots \gamma_i^1 \dots \gamma_p^0$ ($i \in \{1, \dots, p\}$, $p > 0$), $(\pi^j)^1 \gamma_0^2 \gamma_1^0 \dots \gamma_p^0$ ($j \in \{1, \dots, q\}$, $q > 0$) and $(\pi^j)^1 \gamma_0^1 \gamma_1^0 \dots \gamma_i^1 \dots \gamma_p^0$ ($i \in \{1, \dots, p\}$, $j \in \{1, \dots, q\}$, $p > 0$, $q > 0$), read:

$$\partial_{\beta_i} \mathcal{T}_r = 0, \quad (11)$$

$$\partial_{\alpha_0}^2 \mathcal{T}_r - r \partial_{\alpha_0} \mathcal{T}_{r-1} = 0, \quad (12)$$

$$\partial_{\alpha_0 \alpha_i} \mathcal{T}_r - r \partial_{\alpha_i} \mathcal{T}_{r-1} = 0, \quad (13)$$

$$2 \partial_{\alpha_0 \delta_0^j} \mathcal{T}_r - r \partial_{\delta_0^j} \mathcal{T}_{r-1} = 0 \quad (14)$$

resp.

$$\partial_{\alpha_i \delta_0^j} \mathcal{T}_r + \partial_{\alpha_0 \delta_i^j} \mathcal{T}_r - r \partial_{\delta_i^j} \mathcal{T}_{r-1} = 0. \quad (15)$$

These partial equations will allow to compute all equivariant operators.

5 Determination of the equivariant operators

Proposition 1 *Equivariant operators $T \in \mathcal{I}_{p,q}^{k,\ell}$ are mappings*

$$T : \mathcal{D}_p^k \longrightarrow \mathcal{D}_{p-1}^{k+1}, \quad T : \mathcal{D}_p^k \longrightarrow \mathcal{D}_p^k \quad \text{or} \quad T : \mathcal{D}_p^k \longrightarrow \mathcal{D}_{p+1}^{k-1}.$$

Proof. If $T \in \mathcal{I}_{p,p-2}^{k,k+1}$, it follows from (7) and (10) that $\mathcal{T}_r = c_r \alpha_0^r \det(\alpha_i, \beta_i, \delta_i^j)$ ($r \in \{0, \dots, k\}$, $c_r \in \mathbb{R}$) and from (11) that $\mathcal{T}_r = 0$ ($r \in \{0, \dots, k\}$). Hence the result (see remark 1). ■

5.1 Case $q = p + 1$

Proposition 2 *All spaces $\mathcal{I}_{p,p+1}^{k,k-1}$ vanish, except*

- (i) the spaces $\mathcal{I}_{p,p+1}^{1,0}$ with bases defined by equation (18), and
- (ii) the space $\mathcal{I}_{0,1}^{2,1}$ with basis defined by equation (19).

Proof. Equations (7) and (10) show that $\mathcal{T}_r = c_r \alpha_0^{r-1} \det(\delta_i^j)$ ($r \in \{0, \dots, k\}$, $c_r \in \mathbb{R}$). Moreover, equations (12) and (14) yield

$$(r-1)c_r = r c_{r-1} \quad (r \in \{3, \dots, k\}), \quad (16)$$

$$2(r-1)c_r = r c_{r-1} \quad (r \in \{2, \dots, k\}). \quad (17)$$

If $k \geq 3$, (16) and (17) imply that each invariant vanishes. If $k = 2$, (17) confirms that $c_1 = c_2 = c$ ($c \in \mathbb{R}$). If in addition $p > 0$, (15), written for $r = 2$, gives $c = 0$. If $(k, p) = (2, 0)$ or $k = 1$, all local invariants have the form $\mathcal{T}_0 = 0, \mathcal{T}_1 = c\langle Y, \nu \rangle, \mathcal{T}_2 = c\langle Y, \xi \rangle \langle Y, \nu \rangle$ resp. $\mathcal{T}_0 = 0, \mathcal{T}_1 = c \det(\delta_i^j)$.

In [2], P. Lecomte introduced—in a more general framework—some homotopy operator K for the dual d^* of the de Rham differential d .

It's well-known that each $D' \in \mathcal{D}_p^1$ admits a global decomposition $D' = \sum \langle \Lambda, L_X \cdot \rangle + \sum \langle \Omega, \cdot \rangle$, where the sums are finite, where Λ and Ω are anti-symmetric contravariant p -tensors on M , and where X denotes a vector field of M . Similarly, any differential operator $D'' \in \mathcal{D}_0^2$ may be written $D'' = \sum \Lambda L_X \circ L_Y + \sum \Omega L_Z + \sum \Theta$, with $\Lambda, \Omega, \Theta \in N$ and $X, Y, Z \in Vect(M)$.

One easily verifies that

$$K|_{\mathcal{D}_p^1}(D') = \frac{1}{1+p} \sum \langle \Lambda, i_X \cdot \rangle \quad (18)$$

and that

$$K|_{\mathcal{D}_0^2}(D'') = \frac{1}{2} \sum \Lambda (i_X L_Y + L_X i_Y) + \sum \Omega i_Z. \quad (19)$$

For the first formula for instance, it suffices to remember that

$$\langle \Lambda, L_X \omega \rangle \simeq \langle X, \xi \rangle (X_1 \wedge \dots \wedge X_p)(\omega) + (X_1 \wedge \dots \wedge X_p)(\zeta \wedge i_X \omega),$$

with unmistakable notations (see section 2), and to use the local representation of K (see [2]).

These results imply that—in the implicated cases—the homotopy operator is independent of the local coordinates and the partition of unity involved in its construction. Furthermore, the r.h.s. of (18) and (19) is independent of the chosen decomposition of D' and D'' respectively. It's now obvious that

$$K : \mathcal{D}_p^1 \rightarrow \mathcal{D}_{p+1}^0 \text{ and } K : \mathcal{D}_0^2 \rightarrow \mathcal{D}_1^1$$

are equivariant operators.

Finally, the spaces $\mathcal{I}_{p,p+1}^{1,0}$ and $\mathcal{I}_{0,1}^{2,1}$ are generated by K . ■

5.2 Case $q = p - 1$ ($p > 0$)

Proposition 3 *The spaces $\mathcal{I}_{p,p-1}^{k,k+1}$ are one-dimensional vector spaces with basis d^* .*

Proof. Equations (7), (10), and (11) tell us that $\mathcal{T}_r = c_r \alpha_0^r \det(\alpha_i, \delta_i^j)$ ($r \in \{0, \dots, k\}, c_r \in \mathbb{R}$) and equation (13) brings out that $c_r = c$ ($r \in \{0, \dots, k\}, c \in \mathbb{R}$).

Since obviously $d^* \in \mathcal{I}_{p,p-1}^{k,k+1}$, the conclusion follows. ■

5.3 Case $q = p$

Proposition 4 *The spaces $\mathcal{I}_{p,p}^{k,k}$ are one-dimensional, except $\mathcal{I}_{0,0}^{k,k}$ ($k > 0$) and $\mathcal{I}_{p,p}^{1,1}$ ($p > 0$) that have dimension 2. Possible bases are $id, (id, I_0)$ resp. (id, d^*K) , where I_0 is defined by $I_0 : D \rightarrow D(1)id$ (1 stands for the constant function $x \rightarrow 1$).*

Proof. (i) Look first at the case $p = 0$. Equations (7), (10), and (12) show that $\mathcal{T}_0 = c_0$ ($c_0 \in \mathbb{R}$) and $\mathcal{T}_r = c\alpha_0^r$ ($r \in \{1, \dots, k\}, c \in \mathbb{R}$).

If $k = 0$, the space of invariants is generated by id . Otherwise, dimension of $\mathcal{I}_{0,0}^{k,k}$ is 2 and the invariants id and I_0 form a basis.

(ii) If $p > 0$, it follows from equations (7), (8), (10), and (11) that

$$\mathcal{T}_r = c_r \alpha_0^r \det(\delta_i^j) + d_r \alpha_0^{r-1} \sum_{n=1}^p \alpha_n \det_{n0}(\delta_i^j) \quad (r \in \{0, \dots, k\}, c_r, d_r \in \mathbb{R}).$$

In this expression, $\det_{n0}(\delta_i^j)$ denotes the determinant $\det(\delta_i^j)$, where the line $(\delta_n^1, \dots, \delta_n^p)$ has been replaced by $(\delta_0^1, \dots, \delta_0^p)$. When exploiting equations (13) and (14), you find $(r-1)d_r = rd_{r-1}$ ($r \in \{2, \dots, k\}$) resp. $2(r-1)d_r = rd_{r-1}$ ($r \in \{2, \dots, k\}$), so that $d_r = 0$ ($r \in \{0, \dots, k\}$), if $k \geq 2$. Apply now equation (15). If $k \geq 2$, this condition shows that

$$\mathcal{T}_r = c\alpha_0^r \det(\delta_i^j) \quad (r \in \{0, \dots, k\}, c \in \mathbb{R})$$

and if $k = 1$, it entails that $c_1 - c_0 + d_1 = 0$.

If $k = 0$ or $k \geq 2$, invariants are thus generated by id . In the case $k = 1$, dimension is 2 and $id, d^*K \in \mathcal{I}_{p,p}^{1,1}$ are possible generators. ■

Remark 3 Equation (18) reveals that $(d^*K)(D) = (1/(1+p)) \sum \langle \Lambda, i_X d \cdot \rangle$, if $D = \sum \langle \Lambda, L_X \cdot \rangle + \sum \langle \Omega, \cdot \rangle$.

References

- [1] De Wilde M, Lecomte P (1981) Some characterizations of differential operators on vector bundles. In: Butzer P, Feher F (ed) Christoffel, pp 543-549. Basel: Birkhäuser Verlag
- [2] Lecomte P (1994) On some sequence of graded Lie algebras associated to manifolds. Ann Glob Ana Geo **12**: 183-192
- [3] Lecomte P.B.A, Mathonet P, Tousset E (1996) Comparison of some modules of the Lie algebra of vector fields. Indag Math (N.S.) **7**: 461-471
- [4] Peetre J (1959, 1960) Une caractérisation abstraite des opérateurs différentiels. Math Scand **7**: 211-218, **8**: 116-120
- [5] Poncin N (1999) Cohomologie de l'algèbre de Lie des opérateurs différentiels sur une variété, à coefficients dans les fonctions. C R Acad Sci Paris Sér I **328**: 789-794
- [6] Poncin N (2001) On the Cohomology of the Nijenhuis-Richardson Graded Lie Algebra of the Space of Functions of a Manifold. J of Alg **243**: 16-40
- [7] Weyl H (1946) The classical groups, their invariants and representations. Princeton: Princeton Math Series

Université de Luxembourg
Département de Mathématiques
Avenue de la Faïencerie, 162 A
L-1511 Luxembourg, Luxembourg
E-mail: poncin@cu.lu